

Exercises 2

- The following sets are sets of number pairs, i.e., subsets of the cartesian product $\mathbb{R} \times \mathbb{R} = \mathbb{R}^2$:
 $A := \{ (x, y) \in \mathbb{R}^2 \mid y = \frac{2}{3}x - 2 \}$
 $B := \{ (x, y) \in \mathbb{R}^2 \mid y = -|0.5x| + 1 \}$
 $C := \{ (x, y) \in \mathbb{R}^2 \mid x \geq 0 \wedge -x+3 \geq y \geq 0 \}$
 - Visualize each of the sets A, B, C in the cartesian coordinate system. (Make a separate graphical image for each set.)
 - All these sets are relations. Which of them are even functions?
- Let P be the set of all participants of a party and D the set of all drinks which are offered there. Each participant $p_i \in P$ shall get a drink $d_i \in D$ according to his/her preference. In this way, a mapping $f: P \rightarrow D$ shall be defined. What do the following possible properties of f mean in this context?
 - surjectivity
 - injectivity
 - bijectivity
 - Can a participant of the party get two different drinks?
- Let $f: D \rightarrow R$ be the function described by $f(x) = 2x^3 - 1$. The domain D is defined as $D := \{ x \in \mathbb{R} \mid -3 \leq x < 2 \}$. Determine
 - the range $R := f(D)$,
 - a formula for the inverse function f^{-1} .
- Calculate the decimal value of the binary number 1001111.
 - Calculate the hexadecimal representation of the decimal number 999.
 - What is the binary expansion of the value $1/3$?
(Hint: You can do "written division" analogously to the decimal case, but with doubling the remainder in every step instead of multiplying by 10.)
- Let $\vec{a} = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$ and $\vec{b} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$.
 - Draw \vec{a} and \vec{b} in a coordinate system.
 - Determine $2 \cdot \vec{a} + \vec{b}$ by calculation and graphically.
 - Let $\vec{c} = \begin{pmatrix} 1 \\ -11 \end{pmatrix}$. Find a representation of \vec{c} as a linear combination of \vec{a} and \vec{b} .
- Let $\vec{a} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$, $\vec{b} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$. What geometrical objects are described by the following sets?
 - $\{ \vec{b} + t \cdot \vec{a} \mid t \in \mathbb{R} \wedge t \geq 0 \}$
 - $\{ \vec{x} \in \mathbb{R}^2 \mid \vec{a} \cdot \vec{x} = 0 \}$
 - $\{ \vec{x} \in \mathbb{R}^2 \mid \|\vec{x} - \vec{b}\| = 0.5 \}$

7. (a) Are the vectors $\vec{a} = \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$, $\vec{b} = \begin{pmatrix} -3 \\ -2 \\ 5 \end{pmatrix}$ and $\vec{c} = \begin{pmatrix} 4 \\ 1 \\ 0 \end{pmatrix}$ linearly independent?

(b) What is the maximal number of vectors which can be linearly independent in \mathbb{R}^4 ?

8. The points $A = (1; 3)$, $B = (11; 7)$ and $C = (3; 13)$ are given in the cartesian coordinate system.

(a) Let A be the new zero (origin) and calculate the vectors $\vec{b} = \overrightarrow{AB}$ and $\vec{c} = \overrightarrow{AC}$.

(b) Calculate the vector $\vec{d} = \vec{b} + \vec{c} = \overrightarrow{AD}$ and the absolute coordinates of the new point D .

(c) Calculate the inner product $\vec{b} \cdot \vec{c}$ and the angle $\angle(\vec{b}, \vec{c})$.

(d) Extend the vectors by a third dimension (with value 0) and calculate the cross product $\vec{b} \times \vec{c}$.

(e) Calculate the area of the parallelogram spanned by \vec{b} and \vec{c} .